


May is Mental Health Awareness Month and May 1-7, 2023 is **Tardive Dyskinesia Awareness Week**

~600,000
people in the U.S. may
be affected by TD.^{1,2}

Some people living with a mental health condition may also experience **tardive dyskinesia (TD)**, an involuntary movement disorder associated with prolonged use of certain mental health medicines (antipsychotics) that are used to treat bipolar disorder, depression, schizophrenia, and/or schizoaffective disorder.^{1,4-6} Other prescription medicines used to treat upset stomach, nausea, and vomiting may also cause TD. TD is characterized by uncontrollable, abnormal, and repetitive movements of the face, torso, and/or other body parts.⁶

Nearly
70%
of Americans with TD
are undiagnosed.³



Join the Movement!

Over the past five years,
50 states and **Washington, DC**
have declared the first week in May
as TD Awareness Week.

TD can impact patients socially, emotionally, and physically.⁷

According to a survey, patients with diagnosed or suspected TD (n=250) reported the condition moderately or extremely affected them in the following areas^{8*}:



44%

Ability to Work[†]



51%

Ability to Sleep[†]



44%

Leaving the House[†]

*The survey evaluated 51 patients diagnosed with TD and 155 patients who were suspected to have TD, as they experienced involuntary movements.

†Base: Patient ATU 2022: Target patients (n=250). Responses based on survey question: "Since first experiencing involuntary movements, how has your ability to perform the following daily activities been affected, if at all? Rating scale: 1 to 5 when 1 means "not affected at all," and 5 means "extremely negatively affected".

Learn more about TD, living with TD, and how to treat TD by visiting TalkAboutTD.com