

# TD MOD Video

[MUSIC PLAYING]

FEMALE SPEAKER: Monoamines, such as dopamine, are essential for many brain functions, including mood, cognition, and motor control. Healthy dopaminergic neurotransmission involves a careful balance between adequate neurotransmitter release into the synaptic cleft and removal of neurotransmitters from the synapse. Several proteins, including vesicular monoamine transporters or VMAT, reuptake transporters, and presynaptic autoreceptors, are critical to this balance. Dopamine is released from the presynaptic neuron and then binds to receptors located on the presynaptic and/or postsynaptic cell membrane.

Dopamine may also be recycled and along with newly synthesized dopamine, be packaged into synaptic vesicles by vesicular monoamine transporters. Packaged dopamine is then ready for subsequent release. Thereby, promoting synaptic neurotransmission.

Within the nigrostriatal pathway, synaptic neurotransmission involving the monoamine dopamine is critical for sustaining normal motor control. When dopamine release, reuptake, metabolism, and vesicular packaging are balanced, movement is smooth and controlled. However, if the delicate balance of dopaminergic neurotransmission in nigrostriatal areas is disrupted, movement disorders, such as tardive dyskinesia, may result. Symptoms of tardive dyskinesia include involuntary movements of the face, trunk, and extremities.

One way in which dopaminergic neurotransmission may become aberrant in the nigrostriatal region is through prolonged exposure to antipsychotics and other anti-dopaminergic treatments, such as metoclopramide. Antipsychotics aim to reestablish a balance in dopaminergic neurotransmission via antagonism of postsynaptic dopamine D2 receptors in mesolimbic areas. By blocking these mesolimbic dopamine D2 receptors, antipsychotics are often quite effective for ameliorating symptoms of psychosis. However, antipsychotic actions are not confined solely to the mesolimbic area. Chronic antipsychotic-induced lowering of dopaminergic neurotransmission outside of the mesolimbic pathway and particularly in the nigrostriatal area may cause an upregulation or hypersensitivity of dopamine D2 receptors. This dopamine D2 receptor hypersensitivity is hypothesized to underlie the pathophysiology of tardive dyskinesia.